

AD-A185 225


CATION OR SOLVENT-INDUCED SUPERMOLECULAR PHthalocyanine 1/1  
FORMATION: CROWN E..(U) YORK UNIV NORTH YORK (ONTARIO)  
DEPT OF CHEMISTRY N ROBACASHI ET AL. JUN 87  
N00014-84-C-0201

UNCLASSIFIED

F/G 7/3

NL

END  
DATE  
FILMED  
11 87  
DTIC



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

AD-A185 225

OFFICE OF NAVAL RESEARCH

Contract N00014-84-G-0201

Task No. 0051-865

Technical Report #15

12  
DTIC FILE COPY

Cation or Solvent-Induced Supramolecular Phthalocyanine Formation: Crown  
Ether Substituted Phthalocyanines

By

N. Kobayashi and A.B.P. Lever

in

Journal of the American Chemical Society

York University  
Department of Chemistry  
4700 Keele St., North York  
Ontario, Canada M3J 1P3

DTIC  
ELECTE  
S OCT 15 1987 D  
D

Reproduction in whole, or in part, is permitted for any purpose of the United States Government

\*This document has been approved for public release and sale; its distribution is unlimited

\*This statement should also appear in Item 10 of the Document Control Data-DD form 1473. Copies of the form available from cognizant contract administrator

87 10 2 043  
043

## REPORT DOCUMENTATION PAGE

ADA 25 221

1a REPORT SECURITY CLASSIFICATION			1b RESTRICTIVE MARKINGS		
2a SECURITY CLASSIFICATION AUTHORITY Unclassified			3 DISTRIBUTION/AVAILABILITY OF REPORT As it appears on the report		
2b DECLASSIFICATION/DOWNGRADING SCHEDULE					
4 PERFORMING ORGANIZATION REPORT NUMBER(S) Report #15			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a NAME OF PERFORMING ORGANIZATION A.B.P. Lever, York University Chemistry Department		6b OFFICE SYMBOL (If applicable)		7a NAME OF MONITORING ORGANIZATION Office of Naval Research	
6c ADDRESS (City, State, and ZIP Code) 4700 Keele St., North York, Ontario M3J 1P3 Canada		7b ADDRESS (City, State, and ZIP Code) Chemistry Division 800 N. Quincy Street Arlington, VA 22217 U.S.A.			
8a. NAME OF FUNDING/SPONSORING ORGANIZATION		8b OFFICE SYMBOL (If applicable)		9 PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER N00014-84-G-0201	
8c ADDRESS (City, State, and ZIP Code)		10 SOURCE OF FUNDING NUMBERS			
		PROGRAM ELEMENT NO		PROJECT NO	TASK NO
					WORK UNIT ACCESSION NO
11 TITLE (Include Security Classification) Cation or Solvent-Induced Supermolecular Phthalocyanine Formation: Crown Ether Substituted Phthalocyanines					
12 PERSONAL AUTHOR(S) Kobayashi, Nagao and Lever, Alfred B.P.					
13a TYPE OF REPORT Technical		13b. TIME COVERED FROM Aug/86 To Aug/87		14 DATE OF REPORT (Year, Month, Day) June 1987	
15 PAGE COUNT 42					
16 SUPPLEMENTARY NOTATION					
17 COSATI CODES			18 SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP			
19 ABSTRACT (Continue on reverse if necessary and identify by block number) Phthalocyanines with four 15-crown-5 ether voids at the 3,4 positions (MtCRPc) (Mt = H <sub>2</sub> , Zn, Co, Ni and Cu) have been synthesized and characterised. Dimerisation of MtCRPc is induced in solvents such as methanol and by addition of some cations (K <sup>+</sup> , Ca <sup>2+</sup> , and NH <sub>4</sub> <sup>+</sup> ), especially K <sup>+</sup> . Cofacial dimer formation in the presence of these cations proceeds in a two-step three stage process, as indicated by absorption and emission spectroscopy. These species have a highly specific D <sub>4h</sub> eclipsed configuration providing well defined dimeric species for spectroscopic analysis. The ESR spectrum of the cation induced dimeric CuCRPc shows axial symmetry and may be analysed in terms of an inter-planar separation of 4.1 Ångstroms. The <sup>1</sup> H NMR spectra (#19 con'd over)					
20 DISTRIBUTION AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS			21 ABSTRACT SECURITY CLASSIFICATION Unclassified/unlimited		
22a NAME OF RESPONSIBLE INDIVIDUAL Dr. Robert K. Grasselli			22b TELEPHONE (Include Area Code)		22c OFFICE SYMBOL

#19 con'd

of the cation induced metal free and zinc dimers are consistent with an eclipsed configuration. Upper excited state (Soret,  $S_2$ ) emission is observed for the first time in the phthalocyanine series.

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 1113 800 N. Quincy Street Arlington, Virginia 22217-5000	2	Dr. David Young Code 334 NORDA NSTL, Mississippi 39529	1
Dr. Bernard Douda Naval Weapons Support Center Code 50C Crane, Indiana 47522-5050	1	Naval Weapons Center Attn: Dr. Ron Atkins Chemistry Division China Lake, California 93555	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko, Code L52 Port Hueneme, California 93401	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12 high quality	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
DTNSRDC Attn: Dr. H. Singerman Applied Chemistry Division Annapolis, Maryland 21401	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
Dr. William Tolles Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375-5000	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution /	
Availability Codes	
Dist	Avail and/or Special
A1	



ABSTRACTS DISTRIBUTION LIST, 359/627

Dr. Manfred Breiter  
Institut für Technische Elektrochemie  
Technischen Universität Wien  
9 Getreidemarkt, 1160 Wien  
AUSTRIA

Dr. E. Yeager  
Department of Chemistry  
Case Western Reserve University  
Cleveland, Ohio 44106

Dr. R. Sutula  
The Electrochemistry Branch  
Naval Surface Weapons Center  
Silver Spring, Maryland 20910

Dr. R. A. Marcus  
Department of Chemistry  
California Institute of Technology  
Pasadena, California 91125

Dr. J. J. Auborn  
AT&T Bell Laboratories  
600 Mountain Avenue  
Murray Hill, New Jersey 07974

Dr. M. S. Wrighton  
Chemistry Department  
Massachusetts Institute  
of Technology  
Cambridge, Massachusetts 02139

Dr. B. Stanley Pons  
Department of Chemistry  
University of Utah  
Salt Lake City, Utah 84112

Dr. Bernard Spielvogel  
U.S. Army Research Office  
P.O. Box 12211  
Research Triangle Park, NC 27709

Dr. Mel Miles  
Code 3852  
Naval Weapons Center  
China Lake, California 93555

Dr. P. P. Schmidt  
Department of Chemistry  
Oakland University  
Rochester, Michigan 48063

Dr. Roger Belt  
Litton Industries Inc.  
Airtron Division  
Morris Plains, NJ 07950

Dr. Ulrich Stimming  
Department of Chemical Engineering  
Columbia University  
New York, NY 10027

Dr. Royce W. Murray  
Department of Chemistry  
University of North Carolina  
Chapel Hill, North Carolina 27514

Dr. Michael J. Weaver  
Department of Chemistry  
Purdue University  
West Lafayette, Indiana 47907

Dr. R. David Rauh  
EIC Laboratories, Inc.  
Norwood, Massachusetts 02062

Dr. Edward M. Eyring  
Department of Chemistry  
University of Utah  
Salt Lake City, UT 84112

Dr. M. M. Nicholson  
Electronics Research Center  
Rockwell International  
3370 Miraloma Avenue  
Anaheim, California

Dr. Nathan Lewis  
Department of Chemistry  
Stanford University  
Stanford, California 94305

Dr. Hector D. Abruna  
Department of Chemistry  
Cornell University  
Ithaca, New York 14853

Dr. A. B. P. Lever  
Chemistry Department  
~~York University~~  
~~Downsview, Ontario M3J 1P3~~

ABSTRACTS DISTRIBUTION LIST, 359/627

Dr. Martin Fleischmann  
Department of Chemistry  
University of Southampton  
Southampton SO9 5H UNITED KINGDOM

Dr. Alan Bewick  
Department of Chemistry  
The University of Southampton  
Southampton, SO9 5NH UNITED KINGDOM

Dr. John Wilkes  
Department of the Air Force  
United States Air Force Academy  
Colorado Springs, Colorado 80840-6528

Dr. Edward Fletcher  
Department of Mechanical Engineering  
University of Minnesota  
Minneapolis, Minnesota 55455

Dr. R. A. Osteryoung  
Department of Chemistry  
State University of New York  
Buffalo, New York 14214

Dr. Bruce Dunn  
Department of Engineering &  
Applied Science  
University of California  
Los Angeles, California 90024

Dr. Janet Osteryoung  
Department of Chemistry  
State University of New York  
Buffalo, New York 14214

Dr. Elton Cairns  
Energy & Environment Division  
Lawrence Berkeley Laboratory  
University of California  
Berkeley, California 94720

Dr. A. J. Bard  
Department of Chemistry  
University of Texas  
Austin, Texas 78712

Dr. Richard Pollard  
Department of Chemical Engineering  
University of Houston  
Houston, Texas 77004

Dr. Steven Greenbaum  
Department of Physics and Astronomy  
Hunter College  
695 Park Avenue  
New York, New York 10021

Dr. M. Philpott  
IBM Research Division  
Mail Stop K 33/801  
San Jose, California 95130-6099

Dr. Donald Sandstrom  
Boeing Aerospace Co.  
P.O. Box 3999  
Seattle, Washington 98124

Dr. Martha Greenblatt  
Department of Chemistry, P.O. Box 939  
Rutgers University  
Piscataway, New Jersey 08855-0939

Mr. James R. Moden  
Naval Underwater Systems Center  
Code 3632  
Newport, Rhode Island 02840

Dr. Anthony Sammells  
Eltron Research Inc.  
4260 Westbrook Drive, Suite 111  
Aurora, Illinois 60505

Dr. D. Rolison  
Naval Research Laboratory  
Code 6171  
Washington, D.C. 20375-5000

Dr. C. A. Angell  
Department of Chemistry  
Purdue University  
West Lafayette, Indiana 47907

Dr. D. F. Shriver  
Department of Chemistry  
Northwestern University  
Evanston, Illinois 60201

Dr. Thomas Davis  
Polymers Division  
National Bureau of Standards  
Gaithersburg, Maryland 20899

ABSTRACTS DISTRIBUTION LIST, 359/627

Dr. Stanislaw Szpak  
Naval Ocean Systems Center  
Code 633, Bayside  
San Diego, California 95152

Dr. Gregory Farrington  
Department of Materials Science  
and Engineering  
University of Pennsylvania  
Philadelphia, Pennsylvania 19104

Dr. John Fontanella  
Department of Physics  
U.S. Naval Academy  
Annapolis, Maryland 21402-5062

Dr. Micha Tomkiewicz  
Department of Physics  
Brooklyn College  
Brooklyn, New York 11210

Dr. Lesser Blum  
Department of Physics  
University of Puerto Rico  
Rio Piedras, Puerto Rico 00931

Dr. Joseph Gordon, II  
IBM Corporation  
5600 Cottle Road  
San Jose, California 95193

Dr. Joel Harris  
Department of Chemistry  
University of Utah  
Salt Lake City, Utah 84112

Dr. J. O. Thomas  
University of Uppsala  
Institute of Chemistry  
Box 531 Baltimore, Maryland 21218  
S-751 21 Uppsala, Sweden

Dr. John Owen  
Department of Chemistry and  
Applied Chemistry  
University of Salford  
Salford M5 4WT UNITED KINGDOM

Dr. O. Stafstudd  
Department of Electrical Engineering  
University of California  
Los Angeles, California 90024

Dr. Boone Owens  
Department of Chemical Engineering  
and Materials Science  
University of Minnesota  
Minneapolis, Minnesota 55455

Dr. Johann A. Joebstl  
USA Mobility Equipment R&D Command  
DRDME-EC  
Fort Belvoir, Virginia 22060

Dr. Albert R. Landgrebe  
U.S. Department of Energy  
M.S. 68025 Forrestal Building  
Washington, D.C. 20595

Dr. J. J. Brophy  
Department of Physics  
University of Utah  
Salt Lake City, Utah 84112

Dr. Charles Martin  
Department of Chemistry  
Texas A&M University  
College Station, Texas 77843

Dr. Milos Novotny  
Department of Chemistry  
Indiana University  
Bloomington, Indiana 47405

Dr. Mark A. McHugh  
Department of Chemical Engineering  
The Johns Hopkins University  
Baltimore, Maryland 21218

Dr. D. E. Irish  
Department of Chemistry  
University of Waterloo  
Waterloo, Ontario, Canada  
N2L 3G1

DL/1113/87/2

ABSTRACTS DISTRIBUTION LIST, 359/627

Dr. Henry S. White  
Department of Chemical Engineering  
and Materials Science  
151 Amundson Hall  
421 Washington Avenue, S.E.  
Minneapolis, Minnesota 55455

Dr. Daniel A. Buttry  
Department of Chemistry  
University of Wyoming  
Laramie, Wyoming 82071

Dr. W. R. Fawcett  
Department of Chemistry  
University of California  
Davis, California 95616

Dr. Peter M. Blonsky  
Eveready Battery Company, Inc.  
25225 Detroit Road, P.O. Box 45035  
Westlake, Ohio 44145

ABSTRACTS DISTRIBUTION LIST, 051A

Dr. M. A. El-Sayed  
Department of Chemistry  
University of California  
Los Angeles, California 90024

Dr. E. R. Bernstein  
Department of Chemistry  
Colorado State University  
Fort Collins, Colorado 80521

Dr. J. R. MacDonald  
Chemistry Division  
Naval Research Laboratory  
Code 6110  
Washington, D.C. 20375-5000

Dr. G. B. Schuster  
Chemistry Department  
University of Illinois  
Urbana, Illinois 61801

Dr. J. B. Halpern  
Department of Chemistry  
Howard University  
Washington, D.C. 20059

Dr. M. S. Wrighton  
Department of Chemistry  
Massachusetts Institute of Technology  
Cambridge, Massachusetts 02139

Dr. W. E. Moerner  
I.B.M. Corporation  
Almaden Research Center  
650 Harry Rd.  
San Jose, California 95120-6099

Dr. A. B. P. Lever  
Department of Chemistry  
York University  
Downsview, Ontario  
CANADA M3J1P3

Dr. George E. Walrafen  
Department of Chemistry  
Howard University  
Washington, D.C. 20059

Jr. Joe Brandelik  
AFWAL/AAD0-1  
Wright Patterson AFB  
Fairborn, Ohio 45433

Dr. Carmen Ortiz  
Consejo Superior de  
Investigaciones Cientificas  
Serrano 121  
Madrid 6, SPAIN

Dr. Kent R. Wilson  
Chemistry Department  
University of California  
La Jolla, California 92093

Dr. G. A. Crosby  
Chemistry Department  
Washington State University  
Pullman, Washington 99164

Dr. Theodore Pavlopoulos  
NOSC  
Code 521  
San Diego, California 91232

Dr. John Cooper  
Code 6173  
Naval Research Laboratory  
Washington, D.C. 20375-5000

Dr. Joseph H. Boyer  
Department of Chemistry  
University of New Orleans  
New Orleans, Louisiana 70148

Dr. Harry D. Gafney  
Department of Chemistry  
Queens College of CUNY  
Flushing, New York 11367-0904

Dept. of Chemistry, York University, 4700 Keele St.,  
North York (Toronto), Ontario, Canada M3J 1P3

Cation or Solvent-Induced Supramolecular Phthalocyanine Formation: Crown  
Ether Substituted Phthalocyanines.

by Nagao Kobayashi<sup>1\*</sup> and A.B.P. Lever<sup>\*</sup>

Abstract

Phthalocyanines with four 15-crown-5 ether voids at the 3,4 positions (MtCRPc) (Mt = H<sub>2</sub>, Zn, Co, Ni and Cu) have been synthesized and characterised. Dimerisation of MtCRPc is induced in solvents such as methanol and by addition of some cations (K<sup>+</sup>, Ca<sup>2+</sup>, and NH<sub>4</sub><sup>+</sup>), especially K<sup>+</sup>. Cofacial dimer formation in the presence of these cations proceeds in a two-step three stage process, as indicated by absorption and emission spectroscopy. These species have a highly specific D<sub>4h</sub> eclipsed configuration providing well defined dimeric species for spectroscopic analysis. The ESR spectrum of the cation induced dimeric CuCRPc shows axial symmetry and may be analysed in terms of an inter-planar separation of 4.1 Ångstroms. The <sup>1</sup>H NMR spectra of the cation induced metal free and zinc dimers are consistent with an eclipsed configuration. Upper excited state (Soret, S<sub>2</sub><sup>1</sup>) emission is observed for the first time in the phthalocyanine series.

### Introduction

Dimerization, often through aggregation of porphyrins<sup>2</sup> and phthalocyanines<sup>3</sup> has been intensively investigated. In general the mechanism of aggregation is poorly understood, the product may be contaminated with monomer and perhaps with higher aggregates, and the relative orientation of the components is obscure. We report here the preparation of phthalocyanines functionalized at the 3,4-positions with four crown ether voids (MtCRPc),<sup>4</sup> together with their cation or solvent induced dimerization.<sup>5</sup> Unambiguous evidence is presented which indicates the step-wise formation of two dimeric species induced by  $K^+$  ions. The first is non-cofacial, while the second is a rigidly cofacial eclipsed  $D_{4h}$  species. This provides the opportunity to study the uv/visible absorption and emission, ESR, and NMR spectra of well defined dimeric species (metal-free, Zn,Cu,Co and Ni) in terms of exciton theory. Upper excited state  $S_2$  emission is observed in phthalocyanine chemistry for the first time. The results are compared with corresponding data for the tetra-crowned porphyrins,<sup>6,7</sup> where the orientation is twisted. We refer to the dimerisation of the monomeric MtCRPc units for ease of discussion while noting that the binuclear products contain one or more main group ions and are therefore not, strictly speaking, dimers of the mononuclear unit.

### Experimental

i) Measurements: Electronic absorption spectra were recorded with a Perkin Elmer-Hitachi model 340 microprocessor spectrometer. Emission and excitation spectra were obtained with a Varian SF330 spectrofluorimeter with appropriate filters to eliminate scattered light. Fluorescence quantum yields were determined by the use of quinine bisulfate in 1N  $H_2SO_4$  ( $\Phi_F = 0.55$  at 296 K),<sup>8a,b</sup> or free base tetraphenylporphyrin in benzene ( $\Phi_F$

= 0.11).<sup>8c</sup> Data were obtained by a comparative calibration method using the same excitation wavelength and absorbance for the crown species and the calibrants, and the same emission energies.

ESR measurements were obtained using a Varian E4 spectrometer, calibrated with DPPH. FTIR spectra were recorded on a Nicolet SX20 spectrometer using KBr discs. <sup>1</sup>H NMR spectra were recorded on a Varian LH360 spectrometer using deuteriochloroform alone, or containing deuteriomethanol or tetramethylsilane as internal standard. The FAB mass spectra were obtained by courtesy of the Nebraska Center for Mass Spectroscopy by the same method as described previously.<sup>9</sup> Solvents, dimethylformamide, 1,2-dichloroethane, diethylether, methanol, ethanol, etc were either distilled or Spectrograde.

#### 11) Synthesis:

##### 2,3-(3',4'-Dibromobenzo)-1,4,7,10,13-pentaoxacyclopentadeca-2-ene.

Following the dibromination of 1,2-dimethoxybenzene<sup>10</sup>, so-called benzo-15-crown-5<sup>11</sup> was reacted with 2 equivalents of bromine in acetic acid, using a small crystal of iodine as initiator. Recrystallisation from ethanol gave colorless plates in 51% yield. Anal.calcd. for C<sub>14</sub>H<sub>18</sub>Br<sub>2</sub>O<sub>5</sub>: C,39.5; H,4.3. Found: C, 39.6; H,4.3%. <sup>1</sup>H NMR (CDCl<sub>3</sub>, internal Me<sub>4</sub>Si) 7.06(2H,s), 4.25-3.70(16H,m). m.p. 80-81°C.

##### 2,3-(3',4'-Dicyanobenzo)-1,4,7,10,13-pentaoxacyclopentadeca-2-ene.

The above dibrominated benzo-15-crown-5 (8.52g, 0.02mol) and CuCN (5.4g, 0.06mol) were refluxed in dry DMF (80mL) for 5h. After rotary evaporation of about 50-60mL of DMF, concentrated ammonia (200mL, 28%) was added, and air bubbled through the solution for 12h. After washing copiously with water, the dry olive-green product was soxhlet extracted with diethyl ether for 3 days. Benzene recrystallisation provided 2.02g (31%) of colorless small needles. Anal. calcd. for C<sub>16</sub>H<sub>18</sub>N<sub>2</sub>O<sub>5</sub> C, 60.4; H,

5.7; N, 8.8. Found: C, 60.1; H, 5.8; N, 8.5%.  $^1\text{H}$  NMR( $\text{CDCl}_3$ ) 7.14(2H,s), 4.35-3.70(16H,m). IR(KBr)  $\nu$  2223( $\text{C}\equiv\text{N}$ )  $\text{cm}^{-1}$ , m.p. 151-2°C.

**Non-metallated ( $\text{H}_2\text{CRPc}$ ) and metallated ( $\text{MtCRPc}$ ) crowned phthalocyanines**

$\text{H}_2\text{CRPc}$  was prepared by refluxing benzo-15-crown-5 dicyanide (1.24g,  $3.9 \times 10^{-3}$  mol) in 2-dimethylaminoethanol (5mL) for 17h while passing ammonia gas through the solution. The precipitate was re-precipitated from diethylether, washed with water, dried and then chromatographed on basic alumina, initially with chloroform as eluant. The initial yellow band, which was discarded, was followed by a blue band which was eluted with ethanol-chloroform, providing 0.468g (38%) of slightly bluish green powder (Anal, see Table I).  $^1\text{H}$  NMR( $\text{CDCl}_3$ ) 8.02(8H,s), 4.7-3.6(64H,m), -3.41(2H,s).

$\text{ZnCRPc}$  was obtained by reaction of  $\text{H}_2\text{CRPc}$  with a large excess of zinc acetate in refluxing DCE/ethanol, for several days in the dark, until the 4-band Q spectrum of  $\text{H}_2\text{CRPc}$  had disappeared. After removal of the solvent, the residue was washed with water and recrystallised from chloroform-ethanol (78% yield). The FAB mass spectrum showed peaks at m/e values of 1339( $\text{M}^+$ ,7), 956(68), 522(22), 460(58), 424(23), 307(46) and 154(100) (relative intensities in parentheses).

Other  $\text{MtCRPcs}$  ( $\text{Mt} = \text{Co(II)}, \text{Fe(II)}, \text{Ni(II)}, \text{Cu(II)}$ ) were prepared by the "nitrile" method.<sup>12</sup> The benzo-15-crown-5 dinitrile and the appropriate metal acetate were refluxed gently in ethylene glycol<sup>10</sup> for several hours. The solution was filtered while hot, and the residue washed with hot ethanol, hot water, and hot ethanol, then chromatographed on basic alumina, resulting in yields between 26 and 51% of the metallated products. Elemental analyses are presented in Table I.

## Results and Discussion

### 1) Cation complexation leading to step-wise cofacial dimer formation.

In chloroform solution, the several MtCRPc species exhibit spectra typical of well behaved mononuclear metallophthalocyanines<sup>12</sup> generally with a single intense  $\pi-\pi^*$  transition in the range 667-708nm with associated higher energy vibrational components, commonly referred to as the Q band, and a second intense and broader  $\pi-\pi^*$  transition at 300-360nm called the Soret<sup>12</sup> (Table II, Fig.1). The metal free species, H<sub>2</sub>CRPc is similar but shows two closely spaced Q bands because of its lower (D<sub>2h</sub>) symmetry. As will be demonstrated below, dimerization occurs in many other solvents or when certain cations are added to this solution. Such dimerization is readily monitored by the dramatic change in absorption, and in particular by a blue shift in both the Soret and Q band absorption.<sup>13</sup>

In the case of porphyrins functionalised at the methine positions with four benzo-15-crown-5<sup>7a</sup>, increasing addition of K<sup>+</sup>, NH<sub>4</sub><sup>+</sup>, and Ba<sup>2+</sup> ions affected the porphyrin Q band absorption by a) a reduction in intensity, b) broadening and c) a red shift of ca 10nm. On the other hand, Na<sup>+</sup>, Mg<sup>2+</sup> and Ca<sup>2+</sup> did not cause any appreciable change in the visible region absorption. Note that the potassium ion is sandwiched between two crown ether units, and does not reside within the ether ring.

The MtCRPc species behave quite differently. Thus K<sup>+</sup>, NH<sub>4</sub><sup>+</sup>, Ca<sup>2+</sup> and Na<sup>+</sup> all alter the spectra appreciably, though not all in the same fashion, while Ba<sup>2+</sup>, Cs<sup>+</sup> and Mg<sup>2+</sup> have little effect. The ionic radii of cations which produce significant changes in the electronic spectra of the MtCRPc species, are equal or smaller than those required in the porphyrin series.<sup>14</sup> In the crown phthalocyanine case, addition of K<sup>+</sup>, NH<sub>4</sub><sup>+</sup> and Ca<sup>2+</sup>

causes a) a reduction in Q band intensity around 660-700nm (the so-called monomer peak) and an increase in peak intensity around 620-640nm, (the so-called dimer peak) i.e. a blue shift of the Q band of ca 30-60nm, b) a blue shift in the Soret (10-20nm) and c) some broadening in the Q band region (Fig.1A, Table II). Such changes are most obvious for  $K^+$  with respect to the amount needed to affect them.

Absorbance intensities in the 660-700 and 620-640nm ranges change markedly with  $K^+$  concentration up to  $[K^+]/[MtCRPc] = ca 2$ . Previous experience with aggregated<sup>3</sup> and binuclear phthalocyanines<sup>13</sup> shows that the loss of low energy Q band absorption, near 660-700nm, of the monomeric species, and the shift to a broad absorption near 620-640nm is consistent with electronic coupling between a pair (or more) of phthalocyanine units. In this case, as will be documented more fully below, dimerisation is the exclusive process.

From the plot (Fig.2) of the intensity changes with  $[K^+]$ , there is evidence for a two-step three stage process. The first, second and third stages occur approximately at  $[K^+]/[MtCRPc]$  values of 0-0.5, 0.5 - 1.5 and over 1.5, respectively. The  $K^+$  induced two-step three stage process is also obvious from the shift in position of the isosbestic points (Fig.1).<sup>15</sup>

Assuming the monomer-dimer equilibrium (eq.1) for the  $K^+$  triggered spectroscopic change:

K



the specific monomer and dimer concentrations were derived using the method of West and Pearce<sup>16</sup>, using as data the changes of intensity at two wavelengths, as a function of  $[K^+]$ . The pure monomer spectrum<sup>17</sup> was assumed to be present in the absence of  $K^+$ . Using this procedure, proof

of formation of the dimer may be obtained by plotting the calculated  $\log[\text{monomer}]$  against calculated  $\log[\text{dimer}]$ .<sup>18</sup> A slope of 2 will be obtained if eqn.(1) is applicable. Formation of higher oligomers would cause the slope to be more steep than 2. Indeed as revealed in Fig.3 there is a region where the slope is 2.0. When  $K^+$  ions are added to a MtCRPc solution, the monomer concentration decreases sharply, and a region with slope = ca 2.0 continues until  $[K^+]/[\text{MtCRPc}] = 0.5$ . After this point, the slope of the plot approaches zero, especially beyond the region  $[K^+]/[\text{MtCRPc}] \geq 1.0$ .

Thus the monomer-dimer conversion proceeds with a very high formation constant ( $K$  for eqn.1 with  $n = 1$ ,  $(6 \pm 2) \times 10^9 \text{ L}^2 \text{ mol}^{-2}$ ) until two MtCRPc units bind ONE cation. Subsequently the spectroscopic changes in the  $[K^+]/[\text{MtCRPc}] \geq 1.0$  region are no longer a monomer to dimer transformation process, but, rather rearrangement of one form of dimer to another. The region  $0.5 \leq [K^+]/[\text{MtCRPc}] \leq 1.5$  can be considered to be the transition stage where the MtCRPc-cation-MtCRPc complex combines with a 2nd and 3rd cation. Considering that the blue shift in the Q and Soret absorption in the dimers can be explained by an exciton interaction<sup>19</sup>, and that the ESR data for CuCRPc in the presence of  $K^+$  or  $Ca^{2+}$  (vide infra) will indicate the presence of two copper atoms in very close proximity, the last stage ( $[K^+]/[\text{MtCRPc}] \geq 1.5$  in Fig.2) can be ascribed to cofacial complex formation. Thus after formation of a non-cofacial MtCRPc-cation-MtCRPc dimeric complex, increasing cation concentration causes a conformational change in the dimer wherein the two phthalocyanine halves encapsulate two or more cations leading to the formation of the cofacial dimer, i.e. a supermolecular phthalocyanine is attained. Probably encapsulation continues until the available sites are saturated, i.e. until two MtCRPc units bind four cations in rigid eclipsed dimer. The inset of Fig.3 shows

the calculated monomer and dimer concentrations of NiCRPc as a function of  $[K^+]/[NiCRPc]$ . Although not shown, similar behaviour is observed with the other MtCRPc species; thus dimerisation is essentially complete (but not cofacial dimerisation) when  $[K^+]/[MtCRPc] > 1.0$ .

The addition of sodium ions did lead to a change in spectrum (Fig.1B) in contradistinction to the porphyrin series.<sup>7a</sup> The band due to the monomer species decreases in intensity, while there is growth in the band due to the dimeric species. Sodium ions are, however, unable to effect the complete conversion to a spectrum, obtained with potassium ions, typical of the cofacial species.

ii) Solvent induced dimerisation.

The intensity of the MtCRPc Q band peak (around 660-700nm) decreases if the solvent is changed from chloroform to methylene chloride, benzene, DMF, dimethyl sulfoxide, toluene, THF, ethyl acetate, methanol etc. In such circumstances the change in spectrum exactly parallels that observed upon  $K^+$  addition. Such spectroscopic changes are also observed in mixed solvents, e.g. Fig.4a shows the effect of mixing methanol into chloroform, upon the CoCRPc spectrum. With increasing ratio of methanol to chloroform, the peak height at 668nm (mainly monomer) decreases but that at 625nm (mainly dimer) increases. For comparison, Fig.4b shows the corresponding data for addition of  $K^+$ . The dimeric spectrum induced by the methanol solvent ( $CHCl_3:MeOH = 2:1$  v/v) is essentially identical to that obtained at a  $[K^+]/[CoCRPc] = 1.6$  ratio which demonstrates that the mixed solvent is giving rise to cofacial dimer formation. Unfortunately the very low solubility of the metallated crown phthalocyanines in dimer-forming solvents (ca.  $1 \times 10^{-5}M$ ) precludes other methods of characterisation. Note that the solubility of the monomers in chloroform is several orders greater.

### 111) Electron Spin Resonance.

Of the several MtCRPc species investigated, CuCRPc is expected to provide the most definitive ESR spectra. Preliminary studies showed that data collection in a chloroform-methanol mixture provided better data than pure chloroform. The spectrum of CuCRPc obtained in the absence of any cations (Fig.5A) is structureless and seems to be that of a somewhat aggregated species.<sup>20</sup>

Addition of an excess of sodium ions led to the development of a well defined structured signal (Fig.5B-a) in frozen solution, basically very similar to previously observed monomeric copper porphyrin and phthalocyanine species, e.g. copper(II) tetraphenylporphyrin.<sup>21</sup> The spectrum is clearly axial and contains both copper hyperfine and nitrogen (from the four coordinating phthalocyanine nitrogen atoms) super-hyperfine structure. The parameters ( $g_{||} = 2.166$ ,  $g_{\perp} = 2.049$ , and  $A_{||}^{Cu} = 228 \times 10^{-4}$ ,  $A_{\perp}^N = 16.7 \times 10^{-4}$ ,  $A_{||}^N = 14.3 \times 10^{-4} \text{ cm}^{-1}$ ) are typical values for a species of this type. As developed further below, the solution giving rise to this spectrum contains about 50% mononuclear species, and 50% of the non-cofacial dimer almost certainly with sodium ions occupying the ether voids in both cases. In the absence of sodium ions, the solution is partially aggregated resulting in broadening of the resonance signals. In the presence of sodium ions, the resulting positive charge keep the copper atoms separated and a clean highly resolved ESR spectrum is observed.

Dramatic changes in the ESR spectrum of CuCRPc occur when  $K^+$  or  $Ca^{2+}$  ions are added (Fig.5C,D). This observation differs from the corresponding Cu(II) crown porphyrin where  $K^+$ , but not  $Ca^{2+}$ , was effective.<sup>7a</sup> Thus with CuCRPc both cations yield two strong perpendicular transitions in the  $g = 2$  region. These transitions do not exhibit nitrogen super-hyperfine coupling but do show the characteristic seven line pattern expected for

two equivalent Cu(II) ions coupled together.<sup>22</sup> In the half-field region ( $\Delta H = 12$ ), seven equally spaced lines (87-88 gauss separation) are observed. These data provide unmistakable evidence for the formation of a symmetric dimeric molecule. The absence of any ESR signal originating from a monomeric CuCRPc molecule, under these conditions, attests to a high formation constant. Moreover the overall band envelope is essentially identical to those of various well defined symmetric binuclear and coupled copper complexes in the literature.<sup>23</sup> The data in Figs. 5C and 5D also exclude any significant proportion of higher aggregate.

The ESR parameters for these spectra are collected in Table III, having been evaluated by standard procedures.<sup>24</sup> Most importantly it is possible to evaluate the Cu-Cu distance from the zero field splitting parameters.<sup>25</sup> In this fashion the calculated Cu-Cu distance is found to be ca. 4.1 Angstroms, in the presence of both  $K^+$  or  $Ca^{2+}$ . This is nearly identical to the distance estimated from CPK molecular models.<sup>26</sup> Note that in the presence of ammonium ions, dimerisation does occur and two strong perpendicular transitions are observed, but the seven line pattern is not resolved.

#### iv) $^1H$ NMR Spectra.

The  $^1H$  NMR spectrum of ZnCRPc is shown in Fig.6, both in the presence and absence of cations. The spectrum of mononuclear ZnCRPc is complex, in the high field region, due to a large number of chemically inequivalent ether fragments, but the signals are well separated by the ring current anisotropy of the phthalocyanine core. Assignments are shown in Fig.6.

The addition of  $CaCl_2$  to a solution of ZnCRPc ( $[Ca^{2+}]/[ZnCRPc]$  ca. 10) increases the complexity of the high field region (Fig.6b) and the signals spread out, especially downfield. These data are consistent with cofacial dimer formation (Fig.7). The singlet of aromatic protons does not

significantly shift but becomes sharper, due, probably to the restricted rotation of the O-CH<sub>2</sub>-CH<sub>2</sub>-O groups or to the greater rigidity imposed by encapsulation of the cations. The negligible shift of the singlet is expected since the 3,4-protons of one phthalocyanine fall in the "black region" of the anisotropic shielding effect of the second ring,<sup>27</sup> assuming the interplanar distance is close to 4.1 Å.<sup>28</sup>

The pyrrole protons of H<sub>2</sub>CRPc shift from -3.41 ppm to -8.09 ppm with cofacial dimer formation consistent with the structure in Fig.7. This large upfield shift indicates an intense diamagnetic ring current interaction in this dimer.<sup>29</sup>

v) Emission spectroscopy.

Fluorescent emission from the lowest  $\pi$ - $\pi^*$  (Q band) state is common in metal free, zinc and other main group phthalocyanines.<sup>30,31</sup> Both mononuclear H<sub>2</sub>CRPc and ZnCRPc exhibit such emission near 700nm. The excitation spectrum of this emission (Fig.8a) parallels that of other mononuclear phthalocyanines<sup>13</sup> except for a rather prominent excitation near 430nm which is also observed in absorption (Table II). This is rather unusual for a phthalocyanine unit and may involve the ether oxygen lone pairs.

Emission from the upper excited  $\pi$ - $\pi^*$  state (Soret), so-called S<sub>2</sub> emission is not uncommon in porphyrin chemistry.<sup>32-34</sup> We have recently noted that several binuclear and tetranuclear phthalocyanines also exhibit upper excited state emission<sup>35</sup> and also find that both metal free and zinc crown phthalocyanine monomer species exhibit emission in chloroform near 430nm (Fig.8). These are the first examples of phthalocyanine upper excited state emission.

The crown S<sub>2</sub> emission is broader than the S<sub>1</sub> emission and the band shape is similar to that observed for tetrabenzporphyrin free base and its

zinc derivative.<sup>32</sup> The excitation spectra of the  $S_2$  emission is similar, but not identical, to the  $S_1$  excitation spectra (Fig.8A, Table II). Assuming that the  $S_2$  emission originates in the same transition which gives rise to the well developed 420nm absorption, then the Stokes shift for  $S_2$  emission is, as in the porphyrin species, very small.

Using the known quantum yields of quinine sulfate<sup>8a,b</sup> (for  $S_2$ ) and free base tetraphenylporphyrin<sup>8c</sup> for calibration, the quantum yields for  $S_1$  and  $S_2$  emission were determined to be, in chloroform, 0.7 and  $1 \times 10^{-2}$  for  $H_2CRPc$ . Some solution instability precluded an accurate determination for  $ZnCrPc$ . The  $S_1$  quantum yield is very similar to those observed earlier for unsubstituted metal free and zinc phthalocyanine species.<sup>30,31</sup>

Addition of  $K^+$  to metal free and zinc crown Pc monomers in chloroform (concentration  $< 5 \times 10^{-6}M$ ) leads to quenching of both the  $S_1$  and  $S_2$  emission, in a parallel fashion (Fig.8b). The concentration of  $K^+$  ion required to quench the fluorescence intensity of a crowned porphyrin is apparently very much less than the concentration of the cation required to reduce the optical density of the porphyrin in the ground state.<sup>7a</sup> In the crowned phthalocyanine system (Fig.9), on the other hand, the quenching behaviour corresponds almost exactly to that in the absorption study (Fig.2). The Stern-Volmer plot for  $S_1$  emission (Fig.9A) is clearly consistent with step-wise encapsulation of the  $K^+$  ions. In the  $0 < [K^+]/[H_2CRPc] < 0.5$  region where the non-cofacial  $H_2CRPc-K^+-H_2CRPc$  species forms (Figs.2,3), quenching is not marked, though it does occur to a small degree. However with further increase in  $[K^+]$ , quenching is greatly enhanced while above  $[K^+]/[H_2CRPc] = ca. 1.5$  the emission intensity become almost constant. The residual emission intensity has an excitation spectrum identical to that of the monomer species and an intensity about 1/700 of the initial intensity. It is due to the small amount of monomer

species in equilibrium with the cofacial species, implying that the cofacial dimer does not emit.<sup>33</sup>

The quenching of the  $S_2$  emission parallels that of the  $S_1$  quenching except that the degree of quenching is much less marked (Fig.9b). There is also some remaining  $S_2$  emission which appears to arise from the cofacial dimer since it is relatively very much more intense than can be reconciled with the equilibrium monomer content (intensity 2/5 of initial value, upon cofacial dimer formation) (Fig.9b). However the excitation spectrum of this residual  $S_2$  emission is the same as in the monomeric species. There remains the suspicion that the  $S_2$  emission does not indeed originate from the phthalocyanine but from an impurity. This possibility appears excluded by the similarity of its excitation spectrum to the Soret absorption, and the similarity of its quenching behaviour to that of  $S_1$ , but, as noted above, there are some inconsistencies.

The addition of  $Na^+$  ion to a chloroform  $H_2CRPc$  solution also quenches, to some degree, both the  $S_1$  and  $S_2$  emission (Fig.9C). However much higher concentrations of  $Na^+$  are needed and it is clearly much less efficient. In the crown porphyrin case,<sup>7a</sup>  $Na^+$  did not perturb emission. As indicated above,  $Na^+$  ion only takes the species to the second non-cofacial dimer stage.

(vi) Absorption spectra (Table II).

a) Mononuclear species and cofacial dimer.

The mononuclear species have absorption spectra in chloroform (Fig.2, Table II) typical of mononuclear  $MtPc$  species<sup>3</sup> though with a prominent shoulder or peak near 420nm which is not so common. Its appearance is reminiscent of the hyper-porphyrin spectra, showing similar additional absorption but the latter arises as a consequence of the central ion

having an  $(np_z)^2$  pair of electrons, e.g.  $Pb(II)$ .<sup>30</sup> This 420nm band shifts but does not disappear upon cofacial dimer formation.

The formation of the cofacial dimeric species provides a convenient method of studying exciton coupling in an eclipsed phthalocyanine species. Some spectra of fully eclipsed cofacial species are shown in Fig. 0, and the data collected in Table II. Note that under the conditions  $[K^+]/[MtCRPc] > 1.5$ , the quantity of mononuclear species in equilibrium with the cofacial dimer is less than 0.2% and therefore its spectrum makes no significant contribution to the spectra reported here for the cofacial species. The most interesting feature of the cofacial dimeric spectra is the single rather broad (compared to the monomer) and symmetrical Q band absorption. Although the molar extinction coefficients of the cofacial dimer Q bands are approximately one-half of the corresponding monomer Q bands, the oscillator strengths are essentially identical, i.e. there is no loss in transition probability in forming the dimer (see Table II).

Thus exciton coupling of the two  $Q S_1$  states leads to two new levels corresponding to an in-phase and out-of-phase coupling. Transitions to the lower, out-of-phase, component are forbidden rigorously in an eclipsed  $D_{4h}$  species.<sup>36</sup> For compounds which show typical monomeric and dimeric spectra ( $MtCRPc$ ,  $M = Zn, Cu, Ni$  and  $Cu$ ), the Q band blue shift upon dimerization is about  $880 - 980 \text{ cm}^{-1}$ . For parallel eclipsed dimers with negligible distortion from  $D_{4h}$  symmetry, the dipole-dipole excitation splitting,  $V$ , is given by eqn.(2):<sup>37</sup>

$$V = e^2 M^2 / 4 \pi \hbar c \epsilon_0 R^3 \quad (2a)$$

$$= 1.16 \times 10^5 M^2 / R^3 \quad (2b)$$

where  $M$ , and  $R$  are the dipole length, and the inter-molecular separation respectively,  $\epsilon_0$  is the permittivity in a vacuum and the other parameters have their usual meaning.  $V$  is given in wavenumbers in (2b) if

M and R are expressed in Angstroms. The dipole length in Angstroms can be estimated from eqn.(3):<sup>30</sup>

$$M^2 = (\epsilon_{\max}/2513G)(\Delta\lambda/\lambda) \quad (3)$$

where G is the degeneracy, 2 in this case,  $\Delta\lambda$  is the halfbandwidth,  $\lambda$  is the wavelength and  $\epsilon_{\max}$  is the molar extinction coefficient of the dimer Q band, respectively. In this fashion the M values are estimated to be about 0.68 - 0.98 Å (Table IV). Alternatively M may be estimated from the oscillator strength<sup>38</sup> yielding the same result. Assuming a distance of R = 4.1 Å, from ESR, the exciton splitting is calculated to be about 770 - 1600 cm<sup>-1</sup>. The exciton splitting may also be estimated as twice the energy difference between the mononuclear Q band and dimeric Q band peak energies, yielding about 1760, 1960, 1910 and 1960 cm<sup>-1</sup> for Ni, Co, Cu and ZnCRPcs respectively. This compares with an exciton splitting of about 2500 cm<sup>-1</sup> for the binuclear bridged cofacial phthalocyanine species.<sup>13</sup> Using these data and eqn.(3), provides an alternate route to estimating inter-metallic distances, leading to the data shown in Table IV; the electronic spectra and electron spin resonance experiments yield comparable results.

The Soret band is much less obviously affected by formation of the cofacial dimer. There is generally a small blue shift but little change in intensity. Nevertheless the shift is comparable to that seen in the Q band.<sup>5a</sup> Thus coupling of the Soret states does appear to have taken place even though little quenching of the S<sub>2</sub> emission apparently occurs.

#### b) Non-cofacial dimer.

The spectra of the non-cofacial dimer, obtained with [K<sup>+</sup>]/[MtCRPc] = 0.5, are collected in Fig.10, while the spectrum of the sodium ion induced non-cofacial dimer of CuCRPc is shown in Fig.1, in both cases in admixture with the mononuclear species. With [K<sup>+</sup>]/[MtCRPc] = 0.5, there is

approximately 50% of monomeric species in equilibrium with the non-cofacial dimer. In Fig.11 the spectra of several non-cofacial dimeric species are shown, obtained by subtracting the spectrum of the mononuclear species from the admixed spectra. One observes that the resulting spectrum is very similar to that of the cofacial species and with an intensity, per dimeric unit, slightly less than that of the cofacial dimer. There is also evidence for the existence of a new shoulder or peak to the blue of the main Q band absorption, and little evidence for any additional absorption to the red of the Q dimer peak. There is no shifting in the Soret band (Fig.1).

The initial non-cofacial dimer, second stage product, reasonably consists of a dimeric molecule whose two halves are joined together at one ether unit by a bridging potassium ion. In fluid solution there may be many conformational forms of this dimeric unit. The electronic spectrum will consist of a 'snap-shot' of all possible conformations. If there is free rotation about the potassium ion, then many conformational forms will have symmetry much lower than  $D_{4h}$ . Such a situation will give rise to substantial absorption to the red of the cofacial dimeric peak,<sup>18,36</sup> yet this does not appear to be observed, rather, additional absorption is seen to the blue. Suppose instead that the ether voids, which are rather bulky, inhibit such rotation, and restrict the molecule to an end-to-end stretched out  $180^\circ$  conformation of  $C_{2h}$  symmetry, viewed edgewise-on as a step conformation. Such a conformation would show two higher energy peaks due to the splitting of the otherwise degenerate exciton coupled state, but transitions to the two lower energy states remain strictly forbidden.<sup>36</sup> Thus the data appear to support such a linear step conformation. Recall, further, that the ESR spectrum of the CuCRPc- $Na^+$ -CuCRPc solution shows no evidence for Cu-Cu coupling,

consistent with the stretched out formulation.

Note further that the two parts of the step must be parallel - if their angle were greater than  $180^\circ$ , i.e. they were tilted away from each other, absorption to the red of the dimer peak would be expected. Such tilting must be absent or small.

It is noteworthy that the mononuclear emission from  $S_1$  is not quenched to any significant extent upon formation of the stage 2 non-cofacial dimer irrespective of whether it is potassium or sodium ion induced. Since the electronic spectra of these species show significant electronic coupling effects, such an observation is rather surprising.

#### Concluding Remarks

The structure of the final stage 3 dimeric species, assigned to a fully eclipsed cofacial  $D_{4h}$  dimer is required when we assume that the two halves are held together by potassium ions bridging crown ether units. With four  $K^+$  ions shared by two MtCRPc units, a parallel arrangement of the Pc planes can surely be anticipated. This structure is supported by the Cu ESR spectrum, by the presence of a single very symmetric, rather weak and rather narrow Q absorption band (compared with the half-bandwidths for more flexible binuclear phthalocyanine species) by the emission and also by the NMR spectra. Note, too, that the cofacial free base  $[H_2CRPc]_2K^+_4$  species also possesses a single relatively narrow Q band absorption suggesting it too possesses  $D_{4h}$  symmetry and that the four hydrogen atoms are, therefore, equally shared between the four pyrrole nitrogen atoms.

This result is contrasted with the cation induced porphyrin dimer<sup>7a</sup>. Due to the position of the 15-crown-5 units (meso position) they were constrained to adopt a lateral complexation of the cation. Consequently,

the macrocyclic rings are rotated by ca.  $45^\circ$  (usually) with respect to each other in a staggered dimer.

This difference in coordination style may explain the difference in cation behaviour in the two series. Thus the larger  $Ba^{2+}$  ion, ineffective in the MtCRPc series, can dimerize the porphyrin whose inter-crown distance can be expanded by further rotation of the porphyrin rings, without weakening the porphyrin-porphyrin  $\pi-\pi$  stabilisation. In the MtCRPc case, expanding the crown-crown distance, will weaken the Pc-Pc interaction.

It is also interesting that  $Ca^{2+}$  which does not dimerise the crown porphyrin and which forms a 1:1 (internal) adduct with 15-crown-5, is, nevertheless, able to generate a cofacial bridged MtCRPc dimer. It is evident that the free energy gained by forming a 1:1 crown ether complex is exceeded, in this case, by forming a 1:2  $Ca^{2+}$ :crown species and the cofacial dimeric  $[MtCRPc]_2[Ca^{2+}]_4$ .

Lastly we would like to stress the importance of the eclipsed cofacial dimers realized in the present study. Although several covalently bound cofacial porphyrin<sup>39</sup> and phthalocyanine<sup>27</sup> dimers have been reported, they are all skewed<sup>30</sup> and/or staggered.<sup>7,27</sup> While not yet proven by x-ray studies, it is likely from the data presented here, that these cofacial crown MtCRPc phthalocyanines are perfectly eclipsed and will therefore be archetypes for future studies in this area.

#### Acknowledgments

We thank the Office of Naval Research (Washington) for financial support. We are also indebted to S.Hirao for help in the synthesis of benzo-15-crown-5 dibromide, and to H.Lam for NMR assistance.

# FIGURE CAPTIONS

- Figure 1. Change of absorption spectra of (A)  $H_2CRPc$  and (B)  $CuCRPc$  (below) by the addition of  $CH_3COOK$  or  $CH_3COONa$  respectively, to 3ml of a  $CHCl_3$  solution of  $H_2CRPc$  or  $CuCRPc$  in a 10mm cell. The  $CH_3COOK$  and  $CH_3COONa$  were each dissolved in  $CHCl_3$ -MeOH(95:5 v/v) and were added using a microsyringe. 60 $\mu$ L were added in all. Arrows indicate the direction of the spectroscopic change. In (B) the final spectrum is that of a  $CH_3COONa$  saturated solution.
- Figure 2. The dependence of absorbance of several  $MtCRPcs$  on  $[K^+]/[MtCRPc]$  for several wavelengths. Experiments were conducted as described in Fig. 1.
- Figure 3. Plots of log [Monomer] versus log [Dimer] for several  $MtCRPc$  in  $CHCl_3$  solution. Experimental data as appeared in Fig. 1 were analyzed using a computer program based on the approximation method of West and Pearce.<sup>16</sup> The solid lines are drawn with the theoretical slope of 2. The inset shows the dependence of monomer and/or dimer concentration of  $NiCRPc$  on  $[K^+]/[NiCRPc]$  calculated for our experimental system.
- Figure 4. (A) Absorption spectra of  $CoCRPc$  in  $CHCl_3$ -MeOH mixtures, a;  $CHCl_3$  alone, b;  $CHCl_3$ :MeOH = 2.9:0.1 v/v, c;  $CHCl_3$ : MeOH = 2.8:0.2 v/v, d;  $CHCl_3$ :MeOH = 2.4:0.6 v/v, e;  $CHCl_3$ :MeOH = 2.0:1.0 v/v. (B) Change in absorption spectrum of  $CoCRPc$  by the addition of  $CH_3COOK$ . Experiments were carried out as described for Fig.1 The spectrum shown by an open arrow was recorded at  $[K^+]/[CoCRPc] = 1.63$ .

Figure 5. ESR spectra of CuCRPc in  $\text{CHCl}_3$  - MeOH (ca. 4:1 v/v), A; in the absence of any cation, and in the presence of B; saturated  $\text{CH}_3\text{COONa}$  at (a) 77 K, and b) room temperature, C;  $\text{CH}_3\text{COOK}$ , ( $[\text{K}^+]/[\text{CuCRPc}] = 4$ ) and D;  $\text{CaCl}_2$  ( $[\text{Ca}^{2+}]/[\text{CuCRPc}] = 10$ )  $[\text{CuCRPc}]_{\text{mM}} = 1$ .

Figure 6. 300 MHz proton NMR spectra of ZnCRPc in A;  $\text{CDCl}_3$  and B;  $\text{CDCl}_3$  containing trace amount of  $\text{CD}_3\text{OD}$  in the presence of  $\text{CaCl}_2$ . Signals marked X and \* are due to solvents.

Figure 7. A proposed structure for the cation-induced dimer of MtCRPc. Solid circles indicate cations such as  $\text{K}^+$  and  $\text{Ca}^{2+}$ .

Figure 8. Emission and excitation spectra of  $\text{H}_2\text{CRPc}$  in A;  $\text{CHCl}_3$  (emission excited at 350nm) B; in the presence of  $\text{CH}_3\text{COOK}$  in  $\text{CHCl}_3$  containing 0.1 v/v percent of MeOH and C; in the presence of  $\text{CH}_3\text{COONa}$  in  $\text{CHCl}_3$  containing 0.1% v/v MeOH.  $[\text{H}_2\text{CRPc}]/\text{M} = 4.91 \times 10^{-6}$ .  $[\text{CH}_3\text{COOK}]/[\text{H}_2\text{CRPc}] = 2.62$ .  $[\text{CH}_3\text{COONa}]/[\text{H}_2\text{CRPc}] = 98$ . Excitation spectra were recorded for emission peaks at A; 704 nm and B; 429nm and C; 429nm.

Figure 9. Stern-Volmer plots of the emission of  $\text{H}_2\text{CRPc}$ . (a)  $\text{S}_1$  emission quenched by  $\text{K}^+$  ion; (b)  $\text{S}_2$  emission quenched with  $\text{K}^+$  ion; (c)  $\text{S}_2$  emission quenched with  $\text{Na}^+$  ion.

Figure 10. Absorption spectra of several MtCRPc species as indicated. Monomeric species (\_\_\_\_)(in chloroform), cofacial  $[\text{MtCRPc}]_2\text{K}^+_4$  species (-----)(in chloroform with 0.1% methanol), and solutions with  $[\text{K}^+]/[\text{MtCRPc}] = 0.5$  containing a roughly 50:50 mixture of monomeric and non-cofacial  $[\text{MtCRPc}]-\text{K}^+-[\text{MtCRPc}]$  species (.....) (in chloroform with 0.02% methanol).

Figure 11. Estimated spectra of several non-cofacial [MtCRPc]-K<sup>+</sup>-[MtCRPc] species obtained from data in Fig.10 by subtraction of the monomeric component. The spectra in the region of the dotted lines are approximate due to uncertainty in the amount of monomeric species to subtract (ca. 40-50%).

## References

- 1) Visiting Professor from the Pharmaceutical Institute, Tohoku University, Sendai, Japan.
- 2) White, W.I. in "The Porphyrins"; Dolphin, D. Ed.; Academic Press: New York, 1978; Vol. V, p.303. Katz, J.J.; Shipman, L.L.; Cotton, T.M.; Janson, T.R. *ibid.* p.401; Kassel, D. Photochem. Photobiol. 1984, 39, 851. Basu, S.; Rohatgi-Mukherjee, K.K.; Arbeloa, I.L. Spectrochem. Acta 1986, 42, 1355. Shelnutt, J.A.; Dobry, M.M.; Satterlee, J.D. J. Phys. Chem. 1984, 88, 4980.
- 3) Sidorov, A.N.; Kotlyar, I.P.; Opt. Spectrosc. 1961, 11, 92. Sheppard, S.E.; Geddes, S.E. J. Am. Chem. Soc. 1944, 66, 1995. Bernauer, K.; Fallab, S. Helv. Chim. Acta 1961, 54, 1287. Kratky, O.; Oelschaeger, H. J. Colloid Interface Sci. 1969, 31, 490. Shelly, Z.A.; Farina, R.D.; Eyring, E.M. J. Phys. Chem. 1970, 74, 617. Shelly, Z.A.; Harward, D.J.; Hemmes, P.; Eyring, E.M. J. Phys. Chem. 1970, 74, 3040. Siegl, H.; Waldmeier, P.; Prijs, B. Inorg. Nucl. Chem. Lett. 1971, 7, 161. Blagrove, R.J.; Gruen, L.C. Aus. J. Chem. 1972, 25, 1972. Monahan, A.R.; Brado, J.A.; DeLuka, A.F. J. Phys. Chem. 1972, 76, 446. 1994. Farina, F.D.; Halko, D.J.; Swinehart, J.H. J. Phys. Chem. 1972, 76, 2343. Boyd, P.D.W.; Smith, T.D. J. Chem. Soc. Dalton Trans. 1972, 839. Abkowitz, M.; Monahan, A.R. J. Chem. Phys. 1973, 58, 2281. Gruen, L.C.; Blagrove, R.J. Aus. J. Chem. 1973, 26, 319. Abel, E.W.; Pratt, J.M.; Whelan, R. J. Chem. Soc. Dalton Trans. 1976, 509. Cookson, D.J.; Smith, T.D.; Boas, J.F. J. Chem. Soc. Dalton Trans. 1976, 1791. Harriman, A.; Richoux, M.C. J. Chem. Soc. Faraday Trans. II 1980, 76, 1618. Darwent, J.R.; McCubbin, I.; Phillips, D. J. Chem. Soc. Faraday Trans. II 1982, 78, 347, 903. Knoesel, R.; Piechocki, C.; Simon, J. J. Photochem. 1985, 29, 445.

- 4) Abbreviations used in this study: Pc, phthalocyaninato dianion; CRPc, tetra(15-crown-5)ed phthalocyaninato dianion; ESR, electron spin resonance; FAB, fast atom bombardment; NMR, nuclear magnetic resonance; DPPH, diphenylpicrylhydrazyl; DCE, 1,2-dichloroethane; DMF, dimethylformamide; THF, tetrahydrofuran; FTIR, fourier transform infrared.
- 5) Preliminary results on CuCRPc have been published: (a) Kobayashi, N.; Nishiyama, Y. J. Chem. Soc. Chem. Commun. 1986, 1462. Two other reports on CuCRPc have appeared independently: (b) Koray, A.R.; Ahsen, V.; Bakaroglu, O. J. Chem. Soc. Chem. Commun. 1986, 932; (c) Hendriks, R.; Sielcken, O.E.; Drenth, W.; Nolte, R.J.M. ibid 1986, 1464.
- 6) Kobayashi, N.; Osa, T. Heterocycles 1981, 15, 675.
- 7) (a) Thanabal, V.; Krishnan, V. J. Am. Chem. Soc 1982, 102, 3463. (b) idem. Inorg. Chem 1982, 21, 3606. (c) idem. ibid 1985, 24, 3253. (d) Chandrashekar, T.K.; Willigen, H.; Ebersole, M.H. J. Phys. Chem. 1985, 89, 3453. Willigen, H.; Chandrashekar, T.K. J. Am. Chem Soc. 1986, 108, 709.
- 8) (a) Melhuish, W.H. J. Phys. Chem. 1960, 64, 762. (b) Demas, J.N.; Crosby, G.A. ibid 1971, 75, 991. (c) Seybold, P.G.; Gouterman, M. J. Mol. Struct. 1969, 31, 1.
- 9) Leznoff, C.C.; Marcuccio, S.M.; Greenberg, S.; Lever, A.B.P.; Tomer, K.B. Can. J. Chem. 1985, 63, 623.
- 10) Metz, J.; Schneider, O.; Hanack, M. Inorg. Chem. 1984, 23, 1065.
- 11) Pedersen, C.J. J. Am. Chem. Soc. 1967, 89, 7017.
- 12) Lever, A.B.P. Adv. Inorg. Chem. Radiochem., 1965, 7, 27.
- 13) Dodsworth, E.S.; Lever, A.B.P.; Seymour, P.; Leznoff, C.C. J. Phys. Chem. 1985, 89, 5698.
- 14) The cavity size of 15-crown-5 is estimated from CPK models to be

1.7-2.2 Å. The ionic diameters of the cations used in this paper are  $Mg^{2+}$  1.60,  $Na^+$  1.90,  $Ca^{2+}$  1.98,  $K^+$  2.66,  $Ba^{2+}$  2.70,  $NH_4^+$  2.84,  $Cs^+$  3.34 Å.

- 15) The shift in the position of the isosbestic points were most clearly observed for the CuCRPc and NiCRPc system. For the CuCRPc system, the isosbestic points of the 1st and 2nd stages appeared at 370 and 641nm, while those for the 2nd and 3rd emerged at ca. 418 and 665nm. For NiCRPc, isosbestic points were similarly observed at 378 and 632nm, and at 404 and 672nm respectively.
- 16) Pearce, S.; West, W. J. Phys. Chem. 1965, 69, 1894.
- 17) Judging from the previously accumulated data,<sup>3</sup> spectra of MtCRPcs in  $CHCl_3$ , are clearly those of monomer at  $[MtCRPc] < 2 \times 10^{-5}M$ .
- 18) Nevin, W.A.; Liu, W.; Lever, A.B.P. Can. J. Chem. 1987 in press.
- 19) Kasha, M.; Rawls, H.R.; Et-Bayoumi, M.A. Pure Appl. Chem. 1965, 11, 371.
- 20) Skorobogaty, A.; Smith, T.D.; Dougherty, G.; Pilbrow, R. J. Chem. Soc. Dalton Trans. 1985, 651 and several refs. cited therein.
- 21) Manoharan, P.T.; Rogers, M.T. in "Electron Spin Resonance of Metal Complexes", Ed. Yen, T.F. Plenum Press, New York, 1979.
- 22) Smith, T.D.; Pilbrow, J.R. Coord. Chem. Rev. 1974, 13, 173. Eaton, S.S.; More, K.M.; Sawant, B.M.; Boymel, P.M.; Eaton, G.R. J. Magn. Reson. 1983, 52, 435. Thomson, C. Q. Rev. Chem. Soc. 1968, 22, 45. Kottis, P.; Lefebure, R. J. Chem. Phys. 1963, 39, 393.
- 23) (a) Eaton, S.S.; More, K.M.; Sawant, B.M.; Eaton, G.R. J. Am. Chem. Soc. 1983, 105, 6560. (b) Eaton, S.S.; Eaton, G.R.; Chang, C.K. ibid 1985, 107, 3117. (c) Chang, C.K. J. Heterocycl. Chem. 1977, 14, 1285. (d) Eaton, S.A.; Eaton, G.R. J. Am. Chem. Soc. 1982, 104, 5002.
- 24) Chikira, M.; Yokoi, H.; Isobe, T. Bull. Chem. Soc. Jpn. 1974, 47

2208. Chikira, M.; Kon, H.; Hawley, R.A.; Smith, K.M. J. Chem. Soc. Dalton Trans. 1979, 246 and some refs. cited therein. Chasteen, N.D.; Belford, R.L. Inorg. Chem 1970, 9, 169.
- 25) It is possible to estimate the Cu-Cu distance from the ratio of the intensity of the forbidden half-field transition ( $\Delta M = \pm 2$ ) to the intensity of the allowed transitions ( $\Delta M = \pm 1$ )<sup>23a,b,d</sup> In our case, this ratio is  $4.1 \times 10^{-3}$ , which corresponds to  $r = 4.15 \text{ \AA}$ .
- 26) The crystal structure of benzo-15-crown-5 reveals that the plane containing ether oxygens makes an angle of  $31^\circ$  with the benzene ring: Hanson, I.R. Acta Crystallogr., Sect B 1978, B34, 1026.
- 27) Janson, T.R.; Kane, A.R.; Sullivan, J.F.; Knox, K.; Kenney, M.E. J. Am. Chem. Soc. 1969, 91, 5210. Kane, A.R.; Yalman, R.G.; Kenney, M.E. Inorg. Chem. 1968, 7, 2558. Esposito, J.N.; Lloyd, J.E.; Kenney, M.E. ibid 1966, 5, 1979.
- 28) Johnson, C.E.; Bovey, F.A. J. Chem. Phys 1958, 29, 1012.
- 29) Satterlee, J.D.; Shelnutt, J.A. J. Phys. Chem. 1964, 68, 5487.
- 30) Gouterman, M. in "The Porphyrins", Dolphin, D. Ed; Academic Press: New York, 1978: Vol. III, p. 1; Gouterman, M. J. Chem. Phys. 1959, 30, 1139; 1960, 33, 1523.
- 31) Vincent, P.S.; Voigt, E.M.; Rieckhoff, K.E. J. Chem Phys. 1971, 55, 4131.
- 32) Bajema, L.; Gouterman, M. J. Mol. Spectrosc. 1971, 39, 421. Zalesski, I.E.; Kotlo, V.N.; Sevchenko, A.N.; Solov'ev, K.N.; Shkirman, S.F. Sov. Phys. Dokl. 1973, 18, 320. idem. ibid 1975, 19, 589. Fielding, P.E.; Mau, A.W.-H. Aust. J. Chem. 1976, 29, 933. Aaviksoo, J.; Freiberg, A.; Savikhin, S.; Stelmakh, G.F.; Tsvirko, M.P. Chem. Phys. Lett. 1984, 111, 275.
- 33) In the porphyrin system, cofacial dimer units emit more weakly than

- monomer: Kagan, N.E.; Mauzerall, D.; Merrifield, R.B. J. Am. Chem. Soc. 1977, 99, 5486. Chang, C.K.; Kuo, M.-S.; Wang, C.-B. J. Heterocycl. Chem. 1977, 14, 943. Chang, C.K. ibid. 1977, 14, 1285.
- Guckel, F.; Schweiser, D.; Collman, J.P.; Bencosme, S.; Evitt, E.; Sessler, J. Chem. Phys. 1984, 86, 161. Mialoco, C.; Giannotti, A.; Maillard, P.; Momeuteau, M. Chem. Phys. Lett. 1984, 112, 87.
- 34) Ohno, O.; Kaisu, Y.; Kobayashi, H. J. Chem. Phys. 1985, 82, 1779. Kaizu, Y.; Maekawa, H.; Kobayashi, H. J. Phys. Chem. 1986, 90, 4234. Selensky, R.; Holten, D.; Windsor, M.W.; Paine III, J.B.; Dolphin, D. Chem. Phys. 1981, 60, 33. Brookfield, R.L.; Ellul, H.; Harriman, A.; Porter, G. J. Chem. Soc. Faraday Trans. II 1986, 82, 219. Kurabayashi, Y.; Kikuchi, H.; Kokubun, H.; Kaizu, Y.; Kobayashi, H. J. Phys. Chem. 1984, 88, 1308.
- 35) Mannivannan, V.; Kobayashi, N.; Nevin, W.A.; Seymour, P.; Lever, A.B.P.; Leznoff, C.C. unpublished data.
- 36) Gouterman, M.; Holten, D.; Lieberman, E. Chem. Phys. 1977, 25, 139.
- 37) Hochstrasser, R.H. "Molecular Aspects of Symmetry," W.A. Benjamin Inc. New York, 1966, p.311; Sharp, J.H.; Lardon, M. J. Phys. Chem. 1968, 72, 3230.
- 38) Lever, A.B.P. Inorganic Electronic Spectroscopy, 2nd Edn., Elsevier Science Publ., Amsterdam, 1984.
- 39) Collman, J.P.; Chong, A.O.; Jameson, B.G.; Oakley, R.T.; Rose, E.; Schmittou, E.R.; Ibers, J.A. J. Am. Chem. Soc., 1981, 103, 516. Filler, J.P.; Ravichandran, K.G.; Abdalmuhdi, A.; Tulinsky, A.; Chang, C.K. ibid. 1986, 108, 417.

Table I: Elemental Analytical Data of MtCRPcs.<sup>a</sup>

Compound	C		H		N	
	Found	Calcd (%)	Found	Calcd (%)	Found	Calcd (%)
H <sub>2</sub> CRPc	59.83	60.27	5.74	5.85	8.78	8.79
ZnCRPc	57.25	57.42	5.37	5.42	8.37	8.37
CuCRPc	57.33	57.50	5.36	5.43	8.29	8.38
NiCRPc	57.51	57.71	5.35	5.45	8.30	8.41
CoCRPc	57.54	57.70	5.38	5.45	8.29	8.41

<sup>a</sup> Theoretical values are calculated for  $C_{64}H_{72}N_8O_{20}Mt_1$  except for H<sub>2</sub>CRPc ( $C_{64}H_{72}N_8O_{20}H_2$ ).

**Table II** Characteristic Absorption Bands of  $\text{MnCRPcs}^a$ , nm( $\epsilon$ )

Compound	Monomeric Species	$f^{b,c}$	Half- Bandwidth $\text{cm}^{-1}$	Dimeric Cofacial Species	$f$	Half- Bandwidth $\text{cm}^{-1}$
$\text{H}_2\text{CRPc}$	700(33400) 662(27500) 645(11800) 601(6200) 421(8100) 347(30800)			639(17500)    402(sh) <sup>d</sup> 333(26400)		
$\text{ZnCRPc}$	677(96700) 610(17300) 420(sh) 352(61200)	0.16	350	635(51000) 580(sh)  342(53800)	0.17	730
$\text{CuCRPc}$	676(129400) 610(31200) 409(23300) 338(57400) 292(54100)	0.29	480	635(74000)  389(21400) 328(51400)	0.35	1050
$\text{NiCRPc}$	667(70000)  638(22300) 603(17700) 401(14900)	0.17	520	630(47000)   385(13700) 360(13400)	0.18	845

CoCRPc	668(59400)	0.20	750	627(41200)	0.26	1400
	608(19800)			571(sh)		
	400(13400)			390(sh)		
	329(sh)					
	297(49300)			300(45900)		

- 
- <sup>a</sup> Monomer spectra were collected in  $\text{CHCl}_3$ , while those of dimer were in  $\text{CHCl}_3$  containing ca 0.1% MeOH and in the presence of  $\text{K}^+(\text{CH}_3\text{COOK})$ .  $[\text{K}^+]/[\text{MtCRPc}]=4$ . Extinction coefficient ( $\epsilon$ ) is per phthalocyanine unit.
- <sup>b</sup> Oscillator strength. <sup>c</sup> Oscillator strengths for the monomeric species are lower limits excluding higher energy vibrational satellites.
- <sup>d</sup> sh means shoulder.

Table III. Magnetic Parameters<sup>24</sup> of Cation-Induced CuCRPc Cofacial Dimers<sup>a</sup>

Compound	$g_1$	$g_2$	$A_1$	$A_h$	$D_1$	$D_2$	Frequency Cu-Cu <sup>b</sup>	
			<hr/> (gauss)				(MHz)	distance/Å <sup>o</sup>
[CuCRPc] <sub>2</sub> (K <sup>+</sup> ) <sub>4</sub>	2.050	2.152	107	103	371	407	9093	4.2
[CuCRPc] <sub>2</sub> (Ca <sup>2+</sup> ) <sub>4</sub>	2.047	2.141	105	104	375	425	9096	4.1

<sup>a</sup> See Fig.5 for definition of parameters.

<sup>b</sup> Calculated via  $r^3 = 1.39 \times 10^4 \times g_2/D_2$ , where D is in gauss and r is in Ångstroms.<sup>24</sup>




Fig 1




Fig 2




Fig. 3




Fig. 4




Fig. 9




Fig 5 (continued)




Fig 6




Fig. 7




Fig. 8




Fig. 9




Fig 10




Fig 11

ED  
8